

StiffeFire ANTSTAR®

GRANULAR INSECTICIDE

ACTIVE CONSTITUENT: 2 g/kg BIFENTHRIN

SureFire ANTSTAR is our new bifenthrin-based granular insecticide which gives excellent control of ants, ticks, fleas & many turf pests in the garden & outside areas around your home.

SureFire ANTSTAR has these key features:


- 1. Convenient, ready-to-use formulation
- 2. Easy to apply shaker-pack
- 3. Odour-free, low risk product
- 4. Rapid knockdown & long residual action
- 5. Remains effective after rain.

note: for external use only!

GRANULAR INSECTICIDE

Surefire ANTSTAR Granular Insecticide, containing 2g/Kg Bifenthrin, is now registered for use in the control of many pests of turf, and for control of ants, fleas & ticks in the external surrounds and grounds of homes and other buildings.

SITUATION	PEST	RATE kg/100 m2	CRITICAL COMMENTS
Turf (For example: lawns, commercial turf farms, parks, recreational areas, golf courses, bowling greens, sports fields)	Lawn Armyworm (Spodoptera mauritia) Sod Webworm (Herpetogramma licarsisalis)	0.6	Broadcast SureFire AntStar Granular Insecticide with suitable application equipment to ensure uniform coverage over the treated area. To ensure optimum control, irrigate the treated area with up to 4 mm of water immediately after application. Inspect treated areas for continuing activity. Reapply as required. Where a rate range is indicated use lower rates under lower insect pressure and higher rates under higher insect pressure.
	Argentine Stem Weevil adults (Listronotus bonariensis)	0.6-1.2	
	African Black Beetle adults (Heteronychus arator)		
	Billbug adults (Senophorus sp)	1,2-1.8	
	Black ants, Coastal Brown Ant, Funnel Ant, Sugar Ant, Stinging Ant only	0.6-1.2 0.6-2.2	area. Apply granules to areas where ants are active. Where possible apply granules directly to the nests.
			The elimination of Funnel ands from a particular site will generally require more than one application. Initial applications should be broadcast over the affected areas. As the initial numbers of active colonies are reduced, application should shift to targeting active mounds. Apply granules directly to the mound and in the area immediately surrounding active mounds (300 mm radius).
External surrounds of buildings and structures	Black ants, Coustal Brown Ant, Funnel Ant, Sugar Ant, Stinging Ant only	0.6-2.2	Apply granules to areas where ants are active. Where possible apply granules directly to the next. Use the low rate for maintenance treatments or to control light intestations. Use the higher rate for heavy intestations and for maximum residual control. Perimeter Application: Where perimeter pest control is being implemented apply SureFire AntStar Granular insecticide in a 2-3 m band around and adjacent to the building or structure. Pad mounted electrical sites: To reduce or prevent ant mound building activity on electrical pads, uniformly distribute 60 to 220 g SureFire AntStar Granular Insecticide per 10 m2 of mounting pad. When existing transformers are completely or partially filled with soil excavated by resident ants, remove any excess fill and level the remaining soil prior to application.
	Fleas (Adults and Larvae) (Ctenocephalides spp.)	1.1-2.2	Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. For optimum results irrigate the treated areas with up to 10 mm of water immediately after application. Use higher application rates when treating areas with dense ground cover or heavy leaf litter.
	Ticks (Ixodes spp.)	1.1-2.2	Applications should be made in spring to control larvae and nymphs that reside in the soil and leaf litter. DO NOT make spot applications. Treat the entire area where exposure to ticks may occur. Use higher application rates when treating areas with dense ground cover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals. Repeat as necessary to maintain adequate control.



SureFire



See your local stockist:

© PCT International Pty Ltd www.pct.au.com