



Surefire[®] Zinc Phosphide is our new mice control bait for use in broadacre crops.

Surefire[®] Zinc Phosphide is manufactured here in Australia from locally grown quality wheat, which is carefully graded, cleaned and sterilized, before being mixed with food-grade oil and zinc phosphide powder.

Surefire[®] Zinc Phosphide works by releasing phosphine gas in the stomach of the mouse immediately it eats the poisoned grain. The gas is rapidly absorbed by the stomach, and it then depresses the central nervous system. Mice die from very small amounts of grain eaten, with more than 90% control being typically achieved within 48hrs of application. Surefire[®] Zinc Phosphide is registered for either aerial or calibrated ground application equipment.

Surefire[®] Zinc Phosphide features:

- 👍 Easily applied- requires no mixing
- 👍 Highly effective, rapid results, low application rates
- 👍 Has low secondary-poisoning risk to non-target animals
- 👍 Degrades in the open, leaving no harmful by-products





Restrains:

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).
 DO NOT apply bait in a trail.
 DO NOT apply bait if heavy rain is imminent.
 DO NOT apply to the outer 50m of crop or within 50m of native vegetation. A 50m buffer zone in crop must be employed to reduce risk to birds and other non target animals.
 DO NOT apply bait unless monitoring of mouse numbers indicate the potential for crop damage of economic significance.

Situation	Pest	Rate kg/ha	Critical Comments
Grain crops, legume crops, canola, safflower, nut crops and pasture. (Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seeds have been sown into crop stubble or other similar ground cover).	Mice	1 kg/ha to achieve an even coverage of 2-3 grains/m ²	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 grains/m ² . It is imperative that all attempts are made to bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals. Surefire Zinc Phosphide Mouse Bait will provide a high level of control in most situations. However in extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess effectiveness of the control operation. For information on the most effective methods of monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.



APVMA Approval № 59797/1107

THIS PRODUCT IS AVAILABLE ONLY TO AUTHORIZED PERSONS LICENSED FOR SCHEDULE 7 POISONS.

THIS PRODUCT IS TOO DANGEROUS TO BE USED IN THE HOME GARDEN.

A WITHHOLDING PERIOD OF 14 DAYS APPLIES FOR CROP HARVESTING OR LIVESTOCK GRAZING OF BAITED AREAS.

AVAILABLE IN 10 & 125Kg pack sizes.



See your local stockist: